Geographic Data Science -Lecture VI

Exploring Space in Data

Dani Arribas-Bel

Today

- ESDA
- Spatial Autocorrelation
- Measures
 - Global
 - Local

ESDA

Exploratory

Spatial

Data

Analysis

[Exploratory]

Focus on discovery and assumption-free investigation

[Spatial]

Patterns and processes that put *space* and *geography* at the core

[Data Analysis]

Statistical techniques

Questions that ESDA helps...

Answer

- Is the variable I'm looking at concentrated over space? Do similar values tend to locate closeby?
- Can I identify any particular areas where certain values are clustered?

Ask

- What is behind this pattern? What could be generating the process?
- Why do we observe certain clusters over space?

Spatial Autocorrelation

Everything is related to everything else, but near things are more related than distant things

Waldo Tobler (1970)

Spatial Autocorrelation

- -Statistical representation of Tobler's law
- -Spatial counterpart of traditional correlation

Degree to which similar values are located in similar locations

Two flavors:

- Positive: similar values \rightarrow similar location (*closeby*)
- Negative: similar values → disimilar location (further apart)

Examples

Positive SA: income, poverty, vegetation, temperature...

Negative SA: supermarkets, police stations, fire stations, hospitals...

Scales

[Global]

Clustering: do values tend to be close to other (dis)similar values?

[Local]

Clusters: are there any specific parts of a map with an extraordinary concentration of (dis)similar values?

Global Spatial Autocorr.

Global Spatial Autocorr.

"Clustering"

Overall trend where the distribution of values follows a particular pattern over space


[Positive] Similar values close to each other (high-high, low-low)


[Negative] Similar values far from each other (high-low)

How to measure it???

Moran Plot

- Graphical device that displays a variable on the horizontal axis against its spatial lag on the vertical one
- Variable and spatial weights matrix are preferably standardized
- Asssessment of the overall association between a variable in a given location and in its *neighborhood*





[Interactive Demo]

Moran's I

Formal test of global spatial autocorrelation

Statistically identify the presence of clustering in a variable

Slope of the Moran plot

Inference based on how likely it is to obtain a map like observed from a purely random pattern

$$I = rac{N}{\sum_i \sum_j w_{ij}(Z_i)(Z_j)} rac{\sum_i \sum_j w_{ij}(Z_i)(Z_j)}{\sum_i (Z_i)^2}$$

Local Spatial Autocorr.

Local Spatial Autocorr.

"Clusters"

Pockets of spatial instability

Portions of a map where values are correlated in a particularly strong and specific way

[High-High] + SA of high values (hotspots)

[Low-Low] + SA of low values (coldspots)

[High-Low] - SA (spatial outliers)

[Low-High] - SA (spatial outliers)


LISAs


Local Indicators of Spatial Association

Statistical tests for spatial cluster detection → Statistical significance

Compares the observed map with many randomly generated ones to see how likely it is to obtain the observed associations for each location

$$I_i = rac{Z_i}{m_2} \sum_j W_{ij} Z_j \hspace*{0.2cm} ; \hspace*{0.2cm} m_2 = rac{\sum_i Z_i^2}{N}$$





Recapitulation

ESDA is a family of techniques to explore and spatially interrogate data

Main function: characterize spatial autocorrelation, which can be explored:

- Globally (e.g. Moran Plot, Moran's I)
- Locally (e.g. LISAs)



Geographic Data Science'17 - Lecture 6 by Dani Arribas-Bel is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.